

1

ERRATA FOR [AIT07]

2

KOJI AOYAMA

3

- Page 78, Line (–8); Lemmas 2.2 and 2.10; Theorems 3.1, 4.1, 4.2, and 4.3:

4

5

q -uniformly smoothness constant \longrightarrow q -uniform smoothness constant

6

7

- Page 81, Line 14:

8

An operator A of C into E is said to be *accretive* if there exists $j \in J(x - y)$ such that

9

10

$$\langle Ax - Ay, j \rangle \geq 0$$

11

for all $x, y \in C$.

12

\longrightarrow

13

An operator A of C into E is said to be *accretive* if for $x, y \in C$ there exists $j \in J(x - y)$ such that $\langle Ax - Ay, j \rangle \geq \underline{0}$.

14

15

REFERENCES

16

- [AIT07] K. Aoyama, H. Iiduka, and W. Takahashi, *Strong convergence of Halpern's sequence for accretive operators in a Banach space*, Panamer. Math. J. **17** (2007), no. 3, 75–89. MR2335474 (2008e:47125)

17

18

19

(K. Aoyama) AOYAMA MATHEMATICAL LABORATORY, KONAKADAI, INAGE-

20

KU, CHIBA, CHIBA 263-0043, JAPAN

21

Email address: aoyama@bm.skr.jp