

1 **ERRATA FOR [1]**

2 KOJI AOYAMA

- 3 • Page 4, Line 15:

4 If  $\inf_n \beta_n > 0$ , then  $\{S_n\}$  is ...

5  $\longrightarrow$

6 If each  $U_n$  is quasinonexpansive and  $\inf_n \beta_n > 0$ , then  $\{S_n\}$  is ...

- 7 • Page 6, Lines 12–13:

8 Thus Lemma 2.11 implies that  $F = \bigcap_n F(U_n) \neq \emptyset$ ,  $\{U_n\}$  is a strongly quasinonex-  
9 pansive sequence, and  $\hat{F}(\{U_n\}) = \bigcap_n F(U_n)$ .

10  $\longrightarrow$

11 Thus Lemma 2.11 implies that  $F = \bigcap_n F(U_n) \neq \emptyset$ , each  $U_n$  is quasinonexpansive,  
12 and  $\hat{F}(\{U_n\}) = \bigcap_n F(U_n)$ .

13 **REFERENCES**

- 14 [1] K. Aoyama and S. Iemoto, *Parallel methods for quasinonexpansive mappings in a Hilbert space* (2024),  
15 available at [arXiv:2409.03242\[math.FA\]](https://arxiv.org/abs/2409.03242).

16 (K. Aoyama) AOYAMA MATHEMATICAL LABORATORY, KONAKADAI, INAGE-KU, CHIBA, CHIBA 263-  
17 0043, JAPAN

18 *Email address:* [aoyama@bm.skr.jp](mailto:aoyama@bm.skr.jp)